

Subject Index

- Acoustic velocity 83
activation energy of diffusion 106f, 283f, 356f, 463f
activity
-, in Fe-Mg olivine 146f
-, models for mica 45
-, composition relationships
-, -, in hornblende 472f
-, -, in ilmenite-hematite solution 485f
-, -, in magnesiohastite 149f
-, -, in olivine 146f
-, -, in scapolite 221
-, -, ulvöspinel-magnetite solution 485f
AFC (assimilation, fractional crystallization) 385
AFM diagram 421f
albite 1, 115, 169, 296f
-, content in plagioclase 205
alkali
-, basalts 132, 254, 305f, 331f, 396f, 439f
-, feldspars 296f
-, mafic magmas 175f
-, olivine basalts 305f
-, to tholeiite transition 439f
almandine 123
Al₂SiO₅-polymorphs 43f
-, stability field 45, 52
amorphous partial melt 442
amphibole
-, bearing gabbro 385f
-, bearing pyroxenite 177
-, plagioclase pairs 206
amphibolite 196f, 472f
-, granulite transition 318f
analytical error 212f
anatectic melt 393
Andean volcanism 241f
andesite 253f, 385
andesite melt 463f
angle of subduction 242
ankerite 419f
anorthite 170
anorthoclase 441f
anorthosite 223, 454
annite 49
antigorite 100f
aqueous
-, fluids 419f
-, inclusions 322, 453f
Ar retention in biotite 283f
arc
-, basalts 264
-, magmatism 241f
argillite 368f
assimilation 241f, 253, 332, 396, 463f
asthenospheric mantle 315f
-, derived magmas 331f
augite 253f

Back arc volcanism 242f
Barrovian regional metamorphism 419f
basalt
-, alkali 132, 254, 305f, 331f, 396f
-, alkali-olivine 305f
-, crystallization process 129f
-, crystallization trend 132, 253f
-, dyke 401f
-, genesis at low pressure 129f
-, melt 109, 175f, 214, 396f, 463f
-, oxidation state of melt 91, 463f
-, pantellerite suite, Ethiopian Rift 34f
-, primitive high alumina 253f
basic igneous rocks 181f, 212f, 365f, 396f
batch melting 308
batholith 162, 264, 397
benmoreite 331f
bikitaite 73f
biotite, Dora Maira Massif 1f
-, alteration, of 72f, 268
-, carbonate granofels 164
-, coordination of cations 43f
-, K-Ar dating, Santa Rosa mylonite 283f
-, muscovite granofels 164
-, phengite gneiss, Dora Maira Massif 1f
-, Rb-Sr dating, Dora Maira Massif 29
-, zone 419f
blocking temperature
-, biotite (K-Ar) 290
-, zircon (U-Pb) 31
Bowen mechanism 144
breakdown of hornblende 209
bridging/non bridging oxygens 90, 463f
brucite 99
bulk partition coefficients 313f

Ca
-, amphiboles 196f
-, pyroxene minima 39
-, pyroxene, low 181f
calc alkaline differentiation trend 267
calc alkaline magmas 179, 411
calcite 99f, 163f, 356f, 368f
calc-silicates 162f, 196f, 356f
carbonic inclusions 318f, 453f
carbonic metamorphism 318f, 453f
carbon isotopes 162f
cataclasis 286f
cation occupancy in hornblende 477
Cenozoic volcanism 305f
channeled fluid flow 162f
charnockite 318f, 385f, 453f
charnockitic granite 453f
chemical analysis
-, albite, Dora Maira Massif 9
-, amphiboles, Dora Maira Massif 5
-, amphibole, Hawaii (xenoliths) 449
-, biotite, Dora Maira Massif 13
-, biotite, Hawaii (xenoliths) 449
-, biotite, Santa Rosa mylonite 267f
-, chlorites, Dora Maira Massif 4
-, clinopyroxene, Ethiopian Rift 35
-, clinopyroxene, Hawaii (xenoliths) 444, 449
-, clinopyroxene, high grade terranes 237
-, dumortierite, Dora Maira Massif 6
-, dravite, Dora Maira Massif 20
-, ellenbergerite, Dora Maira Massif 5
-, epidote, Dora Maira Massif 13
-, garnet, Dora Maira Massif 3f
-, garnet, high grade terranes 236
-, glaucophane, Dora Maira Massif 9
-, hornblende, Grenville orogen 474f
-, jadeite, Dora Maira Massif 3
-, microcline, Dora Maira Massif 13
-, muscovite, Dora Maira Massif 13
-, olivine, Hawaii (xenoliths) 444, 449
-, orthopyroxene, Hawaii (xenoliths) 444, 449
-, paragonite, Dora Maira Massif 3
-, phengite, Dora Maira Massif 3
-, phlogopite, Dora Maira Massif 9
-, plagioclase, high grade terranes 235
-, pyrope, Dora Maira Massif 3f
-, scapolite, Dora Maira Massif 13
-, scapolite, high grade terranes 235
-, spinell, Hawaii (xenoliths) 444, 449
-, vermiculite, Dora Maira Massif 4
-, talc, Dora Maira Massif 3
-, tourmaline, Dora Maira Massif 6
chlorite 1f, 119
-, + cookeite, stability of 72f
-, zone 419f
chloritoid 419f
clinopyroxene 34f, 129f, 181f, 196f
closure temperature (K-Ar) 283f
coesite 1f
cogenetic hypothesis 213
compatible elements 396f
compressibility of silicate liquids 82f
CO₂
-, activity in high grade metamorphism 219f
-, dominated fluid phase 60, 167, 219f, 318f
-, H₂O fluid phase 219f, 276f, 356f, 453f
-, homogenization temperature 217f, 318f, 455f
-, inclusions in xenoliths 439f
-, infiltration 234, 272f
-, influx 318f
-, rich fluid inclusions 271f, 318f, 453f
C-O-H system 58, 158f, 166, 231f, 453f
C-O-H-S system 229f
concordia diagram 22f
conductive heating 315
connected network of fluid 369f
contact metamorphism 356f
continental basalt 305f, 331f, 385
conserved elements 213f
cookeite 72f
-, enthalpy 77f
-, entropy 77f
-, heat capacity 77f
-, + quartz 78f
-, + bikitaite 78f
cooling magma 106f, 283f
cooling age 29f
cooling history 364f
cooling rate 113f, 364f
cordierite stability 55f
-, with H₂O-CO₂ fluid phase 61
correlation trend of elements 212f
critical
-, radius 108f, 120f
-, undercooling 114

- Cr 446f
 Cr²⁺ in spinels from kimberlite 160
 crustal
 -, contamination 243, 331f, 391f
 -, contribution to magma genesis 241f, 331f
 -, shear zones 283
 -, thickening 247f
 cryptoperthite 299f
 crystal structure
 -, clinopyroxene (M_1 , M_2 , T-sites) 34ff
 -, C2/c structure 34ff
 crystal
 -, growth 108ff, 118ff, 483
 -, size distribution (CSD) 108f, 118ff
 crystallization
 -, equilibrium 140
 -, experiments 110f
 -, fractional 142, 253, 268
 -, of olivine 144
 -, path 129, 268
 cummingtonite 480
 cumulate 398f, 439f
- Daphnite 423f
 decarbonation reaction 419f
 decompression of granulites 321
 decrepitation modelling of fluid inclusions 324
 deformation of rocks 181ff, 346f, 394
 deformation mechanism 346
 dehydration equilibria 419f
 -, cookeite + quartz 72f
 depletion of crystallizing component 106f
 denominator in pearce diagram 212f
 density of crystals 106f
 devolatilization 182f, 356f, 368, 419f
 Devonian-Adrian orogeny 197
 diamond 154
 -, inclusion 157f
 differentiation 396f
 diffusion 108f, 119, 193, 483ff
 -, Ar in biotite 283f
 -, coefficients 483f
 -, creep 346f
 -, effect of pressure 487
 -, grain boundary 362, 356f
 -, induced grain boundary migration (DIGM) 353f
 -, of magnesium in forsterite 356f
 -, of oxygen in andesite melt 483f
 -, of oxygen in basalt melt 483f
 -, of oxygen in calcite 356f
 -, of oxygen in forsterite 380f
 -, radius 291
 -, volume 346f, 356f
 diffusivity 108f, 283f, 346f, 483f
 dihedral angle 356f, 388f
 diopside 39, 100, 134f, 170
 disequilibrium mineral assemblages 396f
 dislocation creep 346f
 -, climb accommodated 347
 -, recrystallization accommodated 347
 dislocation density 346
 dissolution 346f, 483
 distribution coefficients 305f, 339
 dolomite 99, 164f
 dravite 11f
 ductile deformation 346
- durantierite 11f
 dunite 158, 439f
 dynamic recrystallization 348ff
- Eaionite 49
 eclogite 253
 edenite substitution 196f
 elastic strain energy 346
 ellenbergerite 11f
 -, U-Pb dating, Dora Maira Massif 29
 empirical calibration, mixing properties of mica 43
 enstatite 101, 134f
 enthalpy
 -, of mixing 108f
 -, of fusion 112f
 -, of olivine 146f
 entropy of mixing 494
 epidote 11, 196f
 -, group minerals 196f
 eutectoidal reaction 192
 excess Gibbs energy of solution 485f
 exsolution 189
- Fayalite activity in olivine 146ff
 Fe
 -, coordination in mica 54
 -, distribution in garnet 124
 -, effect on phase relationships in basalt 134
 -, redox equilibrium 82f, 146f
 -, Ti oxides 181ff, 485ff
 -, wustite (IW) buffer 149, 154f, 485f
 feldspars 298f
 Fe/Mg partitioning in
 -, garnet-chlorite 120
 -, garnet-cordierite 61
 -, garnet-biotite 43
 -, garnet-hypersthene 84
 -, garnet-spinel 66
 -, olivine-magnesiowustite 146ff
 -, orthopyroxene-basalt melt 258
 -, solid phases of basalt 130
 Fenner mechanism 144
 FeO content in basalt melt 129ff
 ferric/ferrous partitioning 85
 -, in hornblende 199, 472ff
 fission track dating 285f
 fluid
 -, absence 97, 181f, 271, 356f
 -, buffering 184, 358, 388f
 -, chronology 458f
 -, entrapment 271, 318f
 -, flow 358f, 388ff
 -, -, channelized 182f, 233, 453f
 -, -, hydrothermal 167
 -, -, meteoric-hydrothermal 182f, 298f
 -, -, pervasive 183f, 234, 388f
 -, flux 419f
 -, -, estimation of 435f
 -, immiscibility (CO_2 - H_2O - NaCl) 368ff
 -, inclusion in granite 298ff
 -, inclusion in granulites 233f, 271f, 453f
 -, -, in calcite 388f
 -, -, decrepitation of 273f, 324f
 -, -, in garnet 271ff, 318f
 -, -, homogenization temperature 319, 455f
 -, -, in quartz 272, 318f, 388f, 453f
 -, -, reequilibration of 319
- , infiltration 368f, 419ff
 -, intergranular 193f
 -, magmatic 298f, 380
 -, meteoric 298f, 481
 -, rock interaction 182ff, 232, 298f, 368f, 419ff
 -, -, prograde 168f, 379f
 -, -, retrograde 168f, 379f
 -, saturation 97
 forsterite 100f, 115, 134f, 356ff
 -, activity in olivine 146ff
 -, marble 356ff
 fractional crystallization 331f, 393, 396f, 463
 fractionation 441f
 fractionation of olivine 306f
- Gabbro 385f, 396f, 439f
 garnet 11, 43ff, 118ff, 271ff
 -, bearing gneisses 275
 -, bearing-pyroxenite 439
 -, biotite geothermometer 43f
 -, ilherzolite 312f
 -, megacryst 31f
 -, stability 55f
 geobarometry 14ff
 -, garnet-orthopyroxene 157
 -, ol-opx-sp, experimental calibration 384
 -, oxygen barometry, Fe-Ti oxides 485ff
 -, phengite 17
 -, plagioclase breakdown 43ff, 129f
 geochronology, Santa Rosa mylonite 283f
 geothermometry
 -, clinopyroxene-orthopyroxene 183
 -, Fe-Ti oxides 183, 485ff
 -, two feldspars 277
 -, garnet-orthopyroxene 277
 glaucophane 11f
 gneiss 318f
 grain
 -, boundary 346f, 357, 371f
 -, -, diffusion 352, 356f
 -, growth effects 356f
 -, nucleation/growth 189, 383f
 -, size distribution 357
 -, size reduction 346f
 granite
 -, charnockitic 453f
 -, Hercynian 29, 285f
 -, Moldanubian 30
 -, Miocene (Andes) 396f
 -, Tertiary (Isle of Skye) 298ff
 granitic crust 22
 granodiorite 283f
 granulite
 -, facies rocks 198f, 219f, 271f, 318f, 385f, 453f, 483f
 -, Precambrian 56f
 graphite 453f
 -, diamond transition 17, 154
 -, Raman spectra of 459f
 -, stability field of 480
 greenschist facies 11, 183, 284, 390, 419f
 grossular 123
- HAB (high aluminina basalts) 253ff
 harzburgite 156f, 439

- hastingsitic hornblende 472f
 hawaiite 331f, 439
 heat capacity of olivine 146f
 hematite-ilmenite solid solution 486f
 H₂O activity in hornblende 472ff
 H₂O saturation 58
 H₂O-CO₂ mixture 58, 61, 93ff, 359f, 368ff, 419f
 hornblende 472ff
 -, K-Ar dating 283f
 HFS elements 266, 309
 HRE elements 249, 309, 391f
 hybrid intrusion 393
 hydrofracture 368f
 hydrogen isotopes in hornblende 473f
 hydrothermal alteration 298f, 338
 hydrothermal veins 381
 hypersthene-aillimanite-quartz stability 55f
- Ideal solution in hornblende 482
 ilmenite 154
 incipient charnockite formation 318f
 incompatible elements 331f
 -, enrichment 305f
 infiltration driven metamorphism 419f
 Insubric line 385
 interfacial energy 110, 369f
 interfacial strain energy 181f, 346
 intergranular medium (IGM) 119
 intergranular melt 193
 intergrowth texture 181f
 interstitial melt 176
 inverse trace element modelling 306f
 island arc basalts 265
 isobaric cooling 328, 459
 isochores 459f
- Jadeite 1f
- K-Ar dating 283ff, 439f
 K-feldspar 1f, 170
 kimberlite, South African 154ff
 kinetics, zeroth/first order 361f
 Kinzigite formation 385f
 kyanite 1f, 76
- Laccolith 212
 lamproite 316
 latite 331f
 lherzolite 256f, 175f, 439
 LIL element
 -, enrichment 253f
 -, enriched fluids 314
 -, /HFS element ratio 305f, 337
 limestone 368f
 Li
 -, mica 72ff
 -, aluminosilicates 72f
 limpid feldspar 298f
 liquid-gas interface 469
 liquidus phase 253f
 liquidus temperature 129f
 lithosphere, vertically zoned 180
 lithospheric
 -, mantle 305ff
 -, thinning 331f
 lopolith 181f, 454
 LRE elements 331f, 391f
 -, depletion 253f, 310
 -, /HFS element ratio 243, 249
 luminescence of calcite 168
- Mafic formation (Ivrea Zone) 385f
 mafic
 -, complex 385f, 399f
 -, granulites, India 229
 -, + silicic magma 385f, 396f
 -, xenoliths 229
 magma
 -, mixing 331f, 396f, 463
 -, chamber 396f
 magmatic underplating 251, 394
 magnesiohastingsite 146ff
 -, iron equilibrium 152
 magnesite 101
 magnetite 183
 -, ulvöspinel solid solution 486f
 mantle
 -, crust transition 241
 -, peridotite 264
 -, source region 305f
 -, -, contamination 241
 -, wedge 241f, 253, 332
 -, xenoliths (lherzolite) 175f
 -, xenoliths (pyroxenite) 175f
 marble 356ff, 368f
 -, cake structure 385f
 marialite 220
 MASH-model 241ff, 250f, 385
 material transfer processes 212f
 meionite 219f
 melt/crystal interface 109
 melting experiments 253f
 melting temperature (fluid incl.) 455f
 metacarbonates 93ff, 162ff, 356f
 metapelites 419f
 -, Dora Maira Massif 22f
 -, phase relationships 43ff, 56f, 419ff
 -, high T-experiments 55f
 metamorphism
 -, Alpine 22
 -, amphibolite facies 163, 196ff, 284, 318f, 385f
 -, contact 356f, 380
 -, fluid absent 219f
 -, garnet in high P/T rocks 118ff
 -, granulite facies 196ff, 219f, 271ff, 318ff, 385f, 453f, 481
 -, greenschist facies 163, 284, 380, 419f
 -, Hercynian 386
 -, high pressure glaucophane schist 380
 -, Lepontine 381
 -, low grade 72f, 419f
 -, metacarbonate rocks 93ff, 162ff, 356f, 368f
 -, prograde regional 162ff, 271f, 379f, 419f
 -, retrograde 271f, 379f
 -, ultra high-pressure 1ff
 metasediments 162f
 metasomatised mantle 175f
 meteoric fluid 298f, 461
 Mg-Al celadonite 43ff
 Mg ± 129f, 305f, 445f
 microcline 299
 microthermometry 271f, 322f, 453f
 mineralization
 -, hydrothermal 72f
 -, pegmatitic 72f
 Miocene magmatism 398
- miscibility gap 485f
 mizzonite 220
 modelling of fluid-rock interaction 171, 422f
 -, without fluid infiltration 422f
 -, with constant fluid composition 423f
 -, with time integrated influx of fluid 423f
 modelling metamorphism in metapelites 419f
 monazite, Dora Maira Massif 22
 -, U-Th-Pb dating 29
 monzodiorite 396f
 MORB 254f, 336
 Mössbauer spectroscopy 202f
 mugearite 331f
 multiple saturation 262f
 muscovite, Dora Maira Massif 1f
 -, coordination of cations 43ff
 mylonite 283ff
- Nabbarro-Herring creep 352
 NaCl-H₂O-CO₂ fluids 368f
 Na distribution between hbl-plag 205f
 N₂-bearing fluid inclusions 453f
 Nd isotopes 331f
 nepheline 175f
 network modifying cations 463f
 NNO buffer 90
 non-ideal solution 146f, 208, 494
 norite 181ff, 385f
 normalization schemes for hornblende 476f
 nucleation 108f, 118f, 346
 -, growth model 126
 -, rate 106f
 -, temperature 108f
 -, time 106f
- Ocean island basalts (OIB) 331f
 oceanic lithosphere 241f
 olivine 129f, 146ff, 154f, 181f, 253f
 -, fractionation 144, 213, 305f
 -, solid solution 148ff, 485f
 -, spinel pairs 157f
 -, replacement 188f
 -, zonation 403
 open magmatic system 397f
 orthopyroxene 154f, 185, 253f
 -, solid solution 485f
 Ostwald ripening 109f, 118ff
 oxidation state of mantle 154f, 384
 oxygen barometry 485f
 oxygen fugacity (fO₂) 34f, 132, 146ff, 183, 453f, 473f, 485f
 -, kimberlite 154ff
 -, mantle 147
 -, metapelites, experiments 56
 -, silicate liquids 82ff, 463f
 oxygen isotopes 162ff, 298f, 331f, 385f
 -, thermometry 167
 -, tracer experiments 359
 oxygen thermobarometry (ol-opx-sp) 154f, 384
- Paragonite, Dora Maira Massif 1f
 parental melt 336, 396f, 447f
 pargasite substitution 204
 pargasitic hornblende 472f
 partial molar volume 82f
 partial melting 463

- , of peridotite 264, 305f, 331f, 439f
- partition coefficient 305f
- pentite, Ethiopian Rift 34f
- peak nucleation rate 106f
- pearce element ratio (PER) 212f
- pelitic schists 419f
- pericline 99
- permeability-porosity relationship 356f
- pervasive influx of CO_2 318
- petalite 73f
- petrogenetic grid 93f
- phengite, Dora Maira Massif 1f
- phlogopite 1f, 170
- , amphibole transition 342
- , dehydration equilibria 219f
- , breakdown 315
- phyllonite 296
- pigeonite 38, 134f, 190
- plagioclase 180, 253f, 346f
- , breakdown barometry 43f, 129f
- , zonation 403
- Pliocene volcanism, New Mexico 331f
- porosity 368f
- pressure solution 346f
- primary magma 253, 305f
- protomylonite 283f
- psammite 368f
- pseudo
 - , compounds 103f
 - , points 103f
 - , univariant curves 103f
- pT-path 13
- , reconstruction 13
- , anticlockwise/clockwise 271, 280, 454
- pyrope 1f
- , isograd 16
- , megacrysts 4
- , quartzite 1f
- , coesite 8
- pyrophyllite 73f
- pyroxene
 - , miscibility gap 39
 - , orthomagmatic 183
 - , monoclinic 185
- pyroxenite veins 176
- Quartz 43f, 73f, 119, 366f
 - , coesite transition 15
 - , monzodiorite 403f
- QFM buffer 34f, 58, 82f, 146f, 154f, 183f, 496f
- QFM buffer (quartz, ferrosilite, magnetite) 497f
- , sapphirine stability 55f
- Rayleigh devolatilization 170
- Rb depletion 314
- Rb/Cs ratio 243, 249
- recoil phenomena of ^{40}Ar 289f
- recrystallization 346f, 359, 371f
- REE pattern 266, 306f
- relative oxygen fugacity 491f
- retrograde alteration, Dora Maira Massif 12
- Rb-Sr isotopes
 - , Dora Maira Massif 24
 - , Quaternary volcanism (Andes) 243f
- rhomboidal oxides (FeTiO_3 - Fe_2O_3) 485f
- rifting 332
- RTF (refilled, trapped, fractionated) 385
- rutile 1f
- Saline fluids 296f, 368f
 - sapphirine 55f
 - , + quartz stability 55f
 - scapolite-bearing
 - , calc-silicates 227
 - , gneisses 227f
 - , granulite gneisses 229
 - , mafic granulites 229
 - , mafic xenoliths 229f
 - , metaorthosites 223f
 - scapolite equilibria 219f
- Schreinemakers method 94f
- shear stress/strain 191
- siderite 419f
- siderophyllite 49
- silica undersaturated melt 175f
- silicate melt 82f, 109
- , structure of 99, 463
- , viscosity of 463f
- silic magmas 485f
- silic volcanic rocks 490f
- simulation of metamorphism 419f
- singular points 94f
- Sm-Nd isotopes
 - , Dora Maira Massif 24
 - , Quaternary volcanism (Andes) 243f
 - , Cenozoic volcanism (California) 305f
- Southern Volcanic Zone (SVZ) 241f
- spinel 134f, 154f, 485f
- , peridotite 154
- spodumene 73f
- solid-fluid interface 372f
- solid-solid interface 372f
- solid solution 119f, 140, 146, 472f
- solution-reprecipitation 356f, 370f
- Sr isotopes 305f, 331f, 385f
- stable isotopes (C, O) 162f, 296f, 331f, 385f
- stepped heating 324
- strain rate 346f
- stratovolcano 241
- sub arc mantle wedge 242f
- subcaldera 439f
- subducted related magmatism 241f, 305f, 331f
- subducted slab 253, 332
- subduction 241f, 253, 314, 331f
- subsolidus
 - , processes 181f
 - , reequilibration 439f
- sulfate scapolite solid solution 220
- sulfur fugacity (f_{S_2}) 229f
- supercooling 181f
- supersaturation 107
- surface energy 357, 368f
- surface excess 373
- surface tension 114
- symmetric solution 146f
- symplectite 181f
- Tak 1f, 99f
- tectonic erosion 241f, 247
- Tertiary extension 332
- textural equilibrium 356f, 368f
- , experiments 356f
- texturally-primary fluid inclusions 272f
- thermal divide 142
- thermally driven metamorphism 419f
- thermobarometry with hornblende 482f
- thermochronology 283f
- thermodynamic models for
 - , rhomboidal Fe-Ti oxides 487f
 - , cubic phases (spinel) 487f
- tholeiite 129f, 179, 254, 331f, 439f
- topotactic relationship 191
- trace element
 - , pattern 331f
 - , ratio 249
- tracer diffusion 463f
- trachyte, 34f, 439f
- transposed brine inclusions 322
- tremolite 100f, 170, 204f
- , activity in hornblende 472f
- tschermakite substitution 196f, 482
- turbid feldspar 296f
- two-phase fluids 374f
- Ultramafite, Ivrea Zone 368
- ultramylonite 283f
- undercooling 106f
- U-Th-Pb isotopes, Dora Maira Massif 24
- Vermiculite 1f
- volatile equilibria 221f
- volume interdiffusion 346f, 361
- Water
 - , activity in high grade metamorphism 228f
 - , saturation 79
 - , rock ratio 163
- websterite 176
- wehrlite 177, 439
- wetting properties of fluids 375
- wollastonite 99
- wustite-magnetite (WM) buffer 157
- Xenoliths 439f
 - , dunite 439f
 - , glasses 175f, 446
 - , syenite 446
 - , two pyroxene gabbro 439f
- Zircon 22
 - , U-Pb dating, Dora Maira Massif 29
- zoisite 170

